

Sepsis Clinical Pathway

Date & time: ____/____/____

Quick SOFA score for sepsis:

The presence of at least 2 of the following is suggestive of severe sepsis/septic shock:

- ☐ Altered mental status GCS < 15 ☐ Respiratory rate ≥ 22 ☐ Systolic blood pressure ≤ 100

Nursing initial evaluation within 5 minutes:

GCS score: ____/15 BP: ____/____/____ mmHg HR: ____ mmHg temperature: ____°C Spo2: ____

Time ordered	Sepsis pathway orders	Time processed						
	<p><u>Investigations:</u></p> <p><input type="checkbox"/> Contact doctor on call</p> <p><input type="checkbox"/> Check vitals (<i>CONNECT MONITOR</i>)</p> <p><input type="checkbox"/> Establish IV access</p> <p><input type="checkbox"/> Insert Foley's catheter</p> <p><input type="checkbox"/> Hourly reported fluid chart (<i>input / output</i>)</p> <p><input type="checkbox"/> Collect CBC, Oncology profile, Coagulation profile, Serum lactate & PCT</p> <p><input type="checkbox"/> Blood Cultures (<i>peripherally / centrally, if app</i>)</p> <p><input type="checkbox"/> Other Cultures: <input type="checkbox"/> Sputum <input type="checkbox"/> Wound <input type="checkbox"/> Urine <input type="checkbox"/> other: _____</p> <p><input type="checkbox"/> VBG & Blood glucose level</p> <p><u>Antibiotics:</u> administer antibiotics within 1 hour or sepsis recognition</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%;"><input type="checkbox"/> Meronem IV 1 gram TDS</td> <td rowspan="4" style="width: 50%; vertical-align: top;"> <u>In case of renal impairment:</u> <input type="checkbox"/> Teicoplanin IV 400 mg BID for 2 days, followed by 400 mg OD or <input type="checkbox"/> Linezolid 600 mg IV BID </td> </tr> <tr> <td><input type="checkbox"/> Tazocin IV 4.5 grams TDS</td> </tr> <tr> <td><input type="checkbox"/> Vancomycin IV 15mg/kg per dose BD</td> </tr> <tr> <td colspan="2"><input type="checkbox"/> Other antibiotic(s): _____</td> </tr> </table> <p><u>IV fluids:</u> Start IV fluids 0.9% NS, 500 ml bolus fast drip over 5 minutes followed by IV infusion at a rate ____ ml/kg (calculated dose of 30 ml/kg in the first hour until the *mean arterial pressure is > 65 mm Hg and/or HR <110) *Mean arterial pressure = (2 x Diastolic) + Systolic / 3</p> <p><u>Other orders:</u> _____ _____ _____</p> <p><u>Inform ICU</u> ICU Doctor's name: _____</p> <p><u>Patient disposition:</u> <input type="checkbox"/> Ward <input type="checkbox"/> MICU <input type="checkbox"/> SICU</p>	<input type="checkbox"/> Meronem IV 1 gram TDS	<u>In case of renal impairment:</u> <input type="checkbox"/> Teicoplanin IV 400 mg BID for 2 days, followed by 400 mg OD or <input type="checkbox"/> Linezolid 600 mg IV BID	<input type="checkbox"/> Tazocin IV 4.5 grams TDS	<input type="checkbox"/> Vancomycin IV 15mg/kg per dose BD	<input type="checkbox"/> Other antibiotic(s): _____		
<input type="checkbox"/> Meronem IV 1 gram TDS	<u>In case of renal impairment:</u> <input type="checkbox"/> Teicoplanin IV 400 mg BID for 2 days, followed by 400 mg OD or <input type="checkbox"/> Linezolid 600 mg IV BID							
<input type="checkbox"/> Tazocin IV 4.5 grams TDS								
<input type="checkbox"/> Vancomycin IV 15mg/kg per dose BD								
<input type="checkbox"/> Other antibiotic(s): _____								

Signature: _____ Date & Time: _____