



مركز الكويت لمكافحة السرطان
Kuwait Cancer Control Center

Exemestane (Aromasin®)



* E X E - 0 0 0 0 - D L - C N - B R *



Ministry of Health

Name:

File #:

Ht (cm):

Nationality:

Civil ID:

Wt (Kg):

Gender/Age:

DOB:

BSA (m²):

Indication(s): HR +ve breast cancer, Neoadjuvant / Adjuvant / Palliative.

Central line: Available NA

Allergies: NKA Yes, specify; _____

Standard Protocol:

DRUG	DOSE	ADMINISTRATION
Exemestane	25 mg PO daily	To be given after meal.
<input type="checkbox"/> Adjuvant: Postmenopausal: Start with Tamoxifen treatment for 2-3 years followed by of Exemestane to complete 5 years.		
<input type="checkbox"/> Palliative: To be continued until disease progression or intolerable toxicity.		

Special instructions: If a potent CYP3A4 inducer (e.g: carbamazepine, Rifampin) is used concomitantly, increase exemestane dosage to 50 mg once daily.

Treatment Description:

Cycle	Date	Exemestane	Physician	Consultant
C# __				
C# __				
C# __				
C# __				
C# __				
C# __				
C# __				
C# __				
C# __				

Important Notes:

Reported grade 3/4 toxicities: None Hematological Non-Hematological

If yes; Did it indicate hospitalization? Yes No

Did it indicate chemo-delay for ≥ 7 days? Yes No

Did it indicate dose reduction? Yes No

Did it indicate G-CSF support? Yes No