



مركز الكويت لمكافحة السرطان
Kuwait Cancer Control Center

Fulvestrant (Faslodex®)



Ministry of Health

Name:

File #:

Ht (cm):

Nationality:

Civil ID:

Wt (Kg):

Gender/Age:

DOB:

BSA (m²):

Indication(s): Metastatic breast cancer in postmenopausal women, HR positive.

Central line: Available NA

Allergies: NKA Yes, specify; _____

Standard Protocol:

DRUG	DOSE	ADMINISTRATION	DAYS
Fulvestrant	500 mg (Loading dose)	Administer IM slowly over 1–2 minutes per injection - one injection into each buttock.	D1, 15, 29
Fulvestrant	500 mg (maintenance)	Administer IM slowly over 1–2 minutes per injection - one injection into each buttock.	D1
To be repeated every 4 weeks until disease progression or intolerable toxicity.			

Treatment Description:

Cycle	Date	Fulvestrant	Physician	Consultant
C# __				
C# __				
C# __				
C# __				
C# __				
C# __				
C# __				
C# __				
C# __				

Important Notes:

Reported grade 3/4 toxicities: None Hematological Non-Hematological

If yes; Did it indicate hospitalization? Yes No

Did it indicate chemo-delay for ≥ 7 days? Yes No

Did it indicate dose reduction? Yes No

Did it indicate G-CSF support? Yes No