



مركز الكويت لمكافحة السرطان
Kuwait Cancer Control Center

RIBOcilib (Kisqali®)



Ministry of Health

Name:

File #:

Ht (cm):

Nationality:

Civil ID:

Wt (Kg):

Gender/Age:

DOB:

BSA (m²):

Indication(s): HR +ve breast cancer, Neoadjuvant / Adjuvant / Palliative.

Central line: Available NA

Allergies: NKA Yes, specify; _____

Standard Protocol:

DRUG	DOSE	ADMINISTRATION	DAYS
RIBOcilib	600 mg PO daily	May take with or without food.	D1 - 21
To be repeated every 4 weeks until disease progression or intolerable toxicity.			

Special instructions: In case of missed dose or vomiting, don't take an additional dose and resume dosing with the next scheduled daily dose.
Avoid concomitant use with strong CYP3A inhibitors and consider an alternative concomitant medication with less potential for CYP3A inhibition.
If a strong CYP3A inhibitor must be coadministered, reduce the RIBOcilib dose to 400 mg/day.
If the strong inhibitor is discontinued, change the RIBOcilib dose (after at least 5 half-lives of the strong CYP3A inhibitor) to the dose used prior to the initiation of the strong CYP3A inhibitor.

Treatment Description:

Cycle	Day	Date	RIBOcilib	Physician	Consultant
C# __	D1 - 21				
C# __	D1 - 21				
C# __	D1 - 21				
C# __	D1 - 21				
C# __	D1 - 21				
C# __	D1 - 21				
C# __	D1 - 21				

Important Notes:

Reported grade 3/4 toxicities: None Hematological Non-Hematological

If yes; Did it indicate hospitalization? Yes No

Did it indicate chemo-delay for ≥ 7 days? Yes No

Did it indicate dose reduction? Yes No

Did it indicate G-CSF support? Yes No