



مركز الكويت لمكافحة السرطان  
Kuwait Cancer Control Center

## BEVAcizumab (7.5 mg/kg every 3 weeks)



Ministry of Health

Name:

File #:

Ht (cm):

Nationality:

Civil ID:

Wt (Kg):

Gender/Age:

DOB:

BSA (m<sup>2</sup>):

Indication(s): Advanced stage colorectal cancer.

Central line:  Available  NA

Allergies:  NKA  Yes, specify; \_\_\_\_\_

Parameters: Initiate treatment only if ANC  $\geq$  1000; HB  $\geq$  80; Plt  $\geq$  100,000; CrCl  $>$  45 ml/min.

Urine dipstick  $\leq$  +2 (If Urine dipstick  $>$  2, Do urine protein/creatinine ratio And give bevacizumab if the ratio  $<$  2).

BP  $\leq$  150/90 mmHg.

### Standard Protocol:

DRUG	DOSE	ADMINISTRATION	DAYS
BEVAcizumab (Avastin®)	7.5 mg/kg	IV in 100 mL NS over 90 min. If the initial infusion is well tolerated, shorten second infusion to 60 min. If the second infusion is well tolerated, shorten the subsequent infusions to 30 min.	D1
<b>To be repeated every 3 weeks until disease progression or intolerable toxicity.</b>			

### Treatment Description:

Cycle	Date	Avastin®	Bl. Pr.	Wt (Kg)	Physician	Consultant
C# __						
C# __						
C# __						
C# __						
C# __						
C# __						
C# __						
C# __						

### Important Notes:

Reported grade 3/4 toxicities:  None  Hematological  Non-Hematological

If yes; Did it indicate hospitalization?  Yes  No

Did it indicate chemo-delay for  $\geq$  7 days?  Yes  No

Did it indicate dose reduction?  Yes  No

Did it indicate G-CSF support?  Yes  No