



مركز الكويت لمكافحة السرطان
Kuwait Cancer Control Center

FLOT (DOCEtaxel / OXALiplatin / 5-Fluorouracil)



Ministry of Health

Name:

File #:

Ht (cm):

Nationality:

Civil ID:

Wt (Kg):

Gender/Age:

DOB:

BSA (m²):

Indication(s): Locally advanced gastric and gastro-oesophageal junction adenocarcinoma.

Central line: Available NA

Allergies: NKA Yes, specify; _____

Parameters: Initiate treatment only If ANC ≥ 1000; HB ≥ 80; Plt ≥ 100, 0; CrCl > 45 ml/min. Baseline ECHO

Preparatory Medications:

Dexamethazone tab 8 mg at 12, 6, and 2 hrs before DOCEtaxel

Pre-treatment Medications: (30-60 min before starting treatment)

Ondansetron 8 mg PO/IV

Dexamethasone 10 mg PO/IV

Chlorphenamine 10 mg PO/IV

Standard Protocol:

DRUG	DOSE	ADMINISTRATION	DAYS
DOCEtaxel	50 mg/m ²	IV in 500 mL NS over 60 min.	D1
OXALiplatin	85 mg/m ²	IV in 500 mL D5W over 60 min.	D1
Leucovorin	200 mg/m ²	IV In 250 mL D5W over 15 min.	D1
5-FU (infusion)	2600 mg/m ²	For outpatient: continuous infusion via 5-FU pump or For inpatient: IV in 1000 mL NS over 22 hr	D1

To be repeated every 2 weeks.
In operable cases: give 4 cycles pre-operatively and another 4 cycles post-operatively.

Special instructions: - The appropriate Dose Band INFUSOR for 5-FU will be applied accordingly.
- Avoid ice chips.

Treatment Description:

Cycle	Date	DOCEtaxel	OXALiplatin	Leucovorin	5-FU (inf.)	Physician	Consultant
C# __							
C# __							
C# __							
C# __							

Important Notes:

Reported grade 3/4 toxicities: None Hematological Non-Hematological

If yes; Did it indicate hospitalization? Yes No

Did it indicate chemo-delay for ≥ 7 days? Yes No

Did it indicate dose reduction? Yes No

Did it indicate G-CSF support? Yes No