



مركز الكويت لمكافحة السرطان
Kuwait Cancer Control Center

IRINotecan / DOCEtaxel



Ministry of Health

Name:

File #:

Ht (cm):

Nationality:

Civil ID:

Wt (Kg):

Gender/Age:

DOB:

BSA (m²):

Indication(s): Unresectable locally advanced or metastatic gastric and gastro-oesophageal junction adenocarcinoma.

Central line: Available NA

Allergies: NKA Yes, specify; _____

Parameters: Initiate treatment only If ANC \geq 1500; HB \geq 80; Plt \geq 100,000; CrCl > 45 ml/min.

Preperatory Medications:

Dexamethazone tab 8 mg at 12, 6, and 2 hrs before DOCEtaxel.

Pre-treatment Medications: (30-60 min before starting treatment)

Ondansetron 8 mg PO/IV
Dexamethasone 10 mg PO/IV
Chlorphenamine 10 mg PO/IV

Standard Protocol:

DRUG	DOSE	ADMINISTRATION	DAYS
IRINotecan	250 mg/m ²	IV in 500 mL D5W over 60 min.	D1
DOCEtaxel	60 mg/m ²	IV In 500 mL NS over 60 min.	D1
To be repeated every 3 weeks until disease progression or intolerable toxicity.			

Treatment Description:

Cycle	Date	IRINotecan	DOCEtaxel	Physician	Consultant
C# __					
C# __					
C# __					
C# __					
C# __					
C# __					

Important Notes:

Reported grade 3/4 toxicities: None Hematological Non-Hematological

If yes; Did it indicate hospitalization? Yes No

Did it indicate chemo-delay for \geq 7 days? Yes No

Did it indicate dose reduction? Yes No

Did it indicate G-CSF support? Yes No