



مركز الكويت لمكافحة السرطان  
Kuwait Cancer Control Center

## Modified XELIRI (AXEPT Trial)



Ministry of Health

Name:

File #:

Ht (cm):

Nationality:

Civil ID:

Wt (Kg):

Gender/Age:

DOB:

BSA (m<sup>2</sup>):

Indication(s): Advanced stage colorectal cancer, palliative.

Central line:  Available  NA

Allergies:  NKA  Yes, specify; \_\_\_\_\_

Parameters: Initiate treatment only if ANC  $\geq$  1500; HB  $\geq$  80; Plt  $\geq$  75,000; CrCl  $>$  45 ml/min.

Pre-treatment Medications: (30-60 min before starting treatment)

Ondansetron 8 mg PO/IV

Dexamethasone 10 mg PO/IV

### Standard Protocol:

DRUG	DOSE	ADMINISTRATION	DAYS
IRINotecan	200 mg/m <sup>2</sup>	IV in 500 mL D5W over 90 min.	D1
CAPEcitabine	800 mg/m <sup>2</sup> PO bid	To be given with a large glass of water within 30 min	D1 - 14

**To be repeated every 3 weeks, usually for 6-8 cycles but can be continued until disease progression or intolerable toxicity.**

### Treatment Description:

Cycle	Day	Date	IRINotecan	CAPEcitabine	Physician	Consultant
C# __	D1			XXXXXXXX		
	D1 - 14		XXXXXXXX			
C# __	D1			XXXXXXXX		
	D1 - 14		XXXXXXXX			
C# __	D1			XXXXXXXX		
	D1 - 14		XXXXXXXX			

### Important Notes:

Reported grade 3/4 toxicities:  None  Hematological  Non-Hematological

If yes; Did it indicate hospitalization?  Yes  No

Did it indicate chemo-delay for  $\geq$  7 days?  Yes  No

Did it indicate dose reduction?  Yes  No

Did it indicate G-CSF support?  Yes  No