



مركز الكويت لمكافحة السرطان  
Kuwait Cancer Control Center

## Trifluridine and tipiracil (Lonsurf®)



\* T R T - 0 0 0 0 0 - D L - C N - G I \*



Ministry of Health

Name:

File #:

Ht (cm):

Nationality:

Civil ID:

Wt (Kg):

Gender/Age:

DOB:

BSA (m<sup>2</sup>):

Indication(s):  Metastatic Colon or rectal carcinoma.

Metastatic gastric and gastro-esophageal carcinoma.

Central line:  Available  NA

Allergies:  NKA  Yes, specify; \_\_\_\_\_

Parameters: Initiate treatment only if ANC  $\geq$  1000; HB  $\geq$  80; Plt  $\geq$  100,000

### Standard Protocol:

DRUG	DOSE	ADMINISTRATION	DAYS
Lonsurf®	35 mg/m <sup>2</sup> PO twice daily	To be given immediately after a meal or up to 12 hrs after.	D1-5 and D8-12
To be repeated every 28 days until disease progression or intolerable toxicity.			

Special instructions: - Round the dose to the nearest 5 mg.

- Don't exceed 80 mg/dose.

- No restrictions with food type.

- if dose missed, don't make up for the missed dose.

### Treatment Description:

Cycle	Day	Date	Lonsurf®	Physician	Consultant
C# __					
C# __					
C# __					
C# __					
C# __					
C# __					

### Important Notes:

Reported grade 3/4 toxicities:  None  Hematological  Non-Hematological

If yes; Did it indicate hospitalization?  Yes  No

Did it indicate chemo-delay for  $\geq$  7 days?  Yes  No

Did it indicate dose reduction?  Yes  No

Did it indicate G-CSF support?  Yes  No