



مركز الكويت لمكافحة السرطان
Kuwait Cancer Control Center

BEP (BLEOmycin / Etoposide / CISplatin)



* B E P - 0 0 0 0 0 - 0 3 - 0 4 - G U *



Ministry of Health

Name:

File #:

Ht (cm):

Nationality:

Civil ID:

Wt (Kg):

Gender/Age:

DOB:

BSA (m²):

Indication(s): Germ-Cell Tumors.

Central line: Available NA

Allergies: NKA Yes, specify; _____

Parameters: Initiate treatment only if ANC ≥ 1000; HB ≥ 80; Plt ≥ 100,000; CrCl > 45 ml/min.

Pre-treatment Medications: (30-60 min before starting treatment)

Akynzeo 1 Capsule PO (300 mg NETUpitant/0.5 mg PALONOssetron) on Day 1

Dexamethasone 10 mg PO/IV

Chlorphenamine 10 mg PO/IV

Standard Protocol:

DRUG	DOSE	ADMINISTRATION	DAYS
Etoposide	100 mg/m ²	IV In 500 mL NS over 60 min.	D1, 2, 3, 4, 5
CISplatin	20 mg/m ²	IV In 500 mL NS over 60 min.	D1, 2, 3, 4, 5
BLEOmycin	30 Units	IV In 50 mL NS over 10 min.	D2, 9, 16

To be repeated every 3 weeks for 2-4 cycles.

Treatment Description:

Pre-hydration: 500 mL NS IV over 30 min on Day 1 - 5.

Cycle	Day	Date	Etoposide	CISplatin	BLEOmycin	
C# __	D1				XXXXXXXX	
	D2					
	D3				XXXXXXXX	
	D4				XXXXXXXX	
	D5				XXXXXXXX	
	D9			XXXXXXXX	XXXXXXXX	
	D16			XXXXXXXX	XXXXXXXX	

Physician (Stamp and signature)

Consultant (Stamp and signature)