PACLItaxel / CARBOplatin / BEVAcizumab







Printed: 13/May/2020

Name: Nationality: Gender/Age:				Civil ID:			Ht (cm): Wt (Kg): BSA (m²):	
		ple malignano ≀ailable □	cies, specify: NA	Allergies	:	Yes, specify; _		
Parameter	Urine If the		2 (If Urine dipst		t ≥ 100,000; CrC e protein/creatine		bevacizumat	
Ondans		8 mg PC	D/IV	starting treatme	ent)			
Standard F	Protocol	:						
DRUG		DOSE		ADMINISTRATION			DAYS	
BEVAcizumab (Avastin®)		10 mg/kg	infusion to	IV in 100 mL NS over 90 min. If the initial infusion is well tolerated, shorten second infusion to 60 min. If the second infusion is well tolerated, shorten the subsequent infusions to 30 min.				
PACLItaxel		80 mg/m²	IV in 500 mL D5W glass bottle over 3 hrs.				D1, 8, 15	
CARBOplatin		AUC 2		IV in 250 mL D5W over 30 min.			D1, 8, 15	
To be see			for Covolog					
io pe rel	peated e	very 4 weeks	s for 6 cycles.					
		ıs: Calvert r	nethod was use		ARBOplatin dos Oplatin dose Is 3		um creatinine	
	struction	s: Calvert r	nethod was use		•		um creatinine.	
Special ins	struction	s: Calvert r	nethod was use		•		um creatinine. Wt (Kg)	
Special ins	struction	s: Calvert r In case c	nethod was use	aximum CARBC	Oplatin dose Is 3	00 mg.		
Special ins	struction Descrip	s: Calvert r In case c	nethod was use	aximum CARBC	Oplatin dose Is 3	00 mg.		
Special ins	struction Descrip	s: Calvert r In case c	nethod was use of AUC 2, the man	aximum CARBC	Oplatin dose Is 3	BI. Pr.	Wt (Kg)	