





Printed: 13/May/2020

Name: Nationality: Gender/Age:		File #: Civil ID: DOB:	Ht (cm): Wt (Kg): BSA (m²):			
Indication(s): Locally advanced Head and Neck Cancer as an induction phase.         Central line:       □ Available       □ NA       Allergies:       □ NKA       □ Yes, specify;						
Parameters: Initiate treatment only If ANC ≥ 1000; HB ≥ 80; Plt ≥ 100,000; CrCl > 45 ml/min.						
Preperatory Medications:  Dexamethazone tab 8 mg at 12, 6, and 2 hrs before DOCEtaxel.						
Pre-treatment Medicate Akynzeo Dexamethasone Chlorphenamine	tions: (30-60 min before sta 1 Capsule PO (300 mg NE 10 mg PO/IV 10 mg PO/IV	,	NOsetron) on Day 1			

## **Standard Protocol:**

DRUG	DOSE	ADMINISTRATION	DAYS	
DOCEtaxel	75 mg/m²	IV In 500 mL NS over 60 min.	D1	
CISplatin	75 mg/m²	IV In 1000 mL NS over 60 min.	D1	
5-Fluorouracil	1000 mg/m²	IV in 1000 mL NS over 22 hr.	D1, 2, 3, 4, 5	
<ul> <li>□ Induction: to be repeated every 3 weeks 3 cycles.</li> <li>□ Palliative: to be repeated every 3 weeks for 4 - 6 cycles, according to patient tolerance.</li> </ul>				

## **Treatment Description:**

Pre-hydration: 1 Liter NS + KCl 20 mEq + MgSO<sub>4</sub> 8 mEq IV over 2 hrs.

Cycle	Day	Date	DOCEtaxel	CISplatin	5-Fluorouracil
C#	D1				
	D2		xxxxxx	xxxxxx	
	D3		xxxxxx	xxxxxx	
	D4		xxxxxx	xxxxxx	
	D5		xxxxxx	xxxxxx	

Post-hydration: 1 Liter NS IV over 2 hrs followed by 150 mL Mannitol 20% IV bolus.				
Physician (Stamp and signature)	Consultant (Stamp and signature)			