ATEZOlizumab (Tecentriq®)







Printed: 13/May/2020

Name: Nationality: Gender/Age:		File #: Civil ID: DOB:		Ht (cm): Wt (Kg): BSA (m²):
Indication(s): ☐ Advanced stage NSCLC. ☐ Extensive Stage SCLC (combined With Etoposide/CARBOplatin). ☐ Urothelial Cacrcinoma. ☐ Triple-Negative Breast Cancer (combined with NAB-PACLitaxel/CARBOplatin). Central line: ☐ Available ☐ NA Allergies: ☐ NKA ☐ Yes, specify;				
Standard	Protocol:			
DRUG		DOSE	ADMINISTRATION	DAYS
ATEZOlizumab		1200 mg IV in 250 mL NS over 60 min. If the initial infusion is well tolerated, shorten the subsequent infusions to 30 min.		horten the
To be re	epeated ev	ery 3 weeks till disease pro	gression or intolerable toxicit	ies.
Treatmer	nt Descript	ion:		
Cycle	Date	ATEZOlizumab	Physician	Consultant
C#				
	ted grade 3		Hematological □ Non-Hemato	blogical
If yes;	Did it indid Did it indid	cate hospitalization? cate chemo-delay for ≥ 7 day: cate dose reduction? cate G-CSF support?	☐ Yes ☐ No s? ☐ Yes ☐ No ☐ Yes ☐ No ☐ Yes ☐ No	