



مركز الكويت لمكافحة السرطان
Kuwait Cancer Control Center

CETUXimab (every week)



* C T X - 0 0 0 0 - 0 1 - C N - M L *



Ministry of Health

Name:

File #:

Ht (cm):

Nationality:

Civil ID:

Wt (Kg):

Gender/Age:

DOB:

BSA (m²):

Indication(s): Recurrent/Advanced stage/Metastatic Colorectal Cancer.

Recurrent/Advanced stage/Metastatic Head And Neck Cancers.

Central line: Available NA

Allergies: NKA Yes, specify; _____

Parameters: Initiate treatment only if ANC \geq 1500; HB \geq 80; Plt \geq 75,000; CrCl > 45 ml/min

Standard Protocol:

DRUG	DOSE	ADMINISTRATION	DAYS
CETUXimab	400 mg/m ² (Loading dose)	IV undiluted over 2 hrs. Then, flush the IV line with 50 mL NS at end of infusion.	1st dose only
CETUXimab	250 mg/m ² (maintenance dose)	IV undiluted over 60 min. Then, flush the IV line with 50 mL NS at end of infusion.	D1
To be repeated every 1 week until disease progression or intolerable toxicity.			

Special instructions: Keep the patient For 1 hr observation period after completion of the 1st And 2nd cycle of CETUXimab infusion.

Treatment Description:

Cycle	Date	CETUXimab	Physician	Consultant
C# __				
C# __				
C# __				
C# __				
C# __				
C# __				

Important Notes:

Reported grade 3/4 toxicities: None Hematological Non-Hematological

If yes; Did it indicate hospitalization? Yes No

Did it indicate chemo-delay for \geq 7 days? Yes No

Did it indicate dose reduction? Yes No

Did it indicate G-CSF support? Yes No