



مركز الكويت لمكافحة السرطان
Kuwait Cancer Control Center

Everolimus (Afinitor®)



Ministry of Health

Name:

File #:

Ht (cm):

Nationality:

Civil ID:

Wt (Kg):

Gender/Age:

DOB:

BSA (m²):

Indication(s): Advanced Neuroendocrinal tumors.

Metastatic breast cancer, HR +ve.

Central line: Available NA

Allergies: NKA Yes, specify; _____

Standard Protocol:

DRUG	DOSE	ADMINISTRATION
Everolimus	10 mg PO daily	To be taken at the same time every day, with or without food.
To be given continuously until disease progression or intolerable toxicity.		

Special instructions:

- Everolimus missed doses may be taken up to 6 hours after regularly scheduled time; if > 6 hours, resume at the next regularly scheduled time.
- Patients taking concomitant ACE inhibitor therapy may be at increased risk for angioedema.
- Avoid concomitant use with: Grapefruit or grapefruit juice, and CLARithromycin.
- Use concomitantly with caution with: Anticonvulsants, Fluconazole, voriconazole, with CCBs (diltiazem, verapamil), and erythromycin.

Treatment Description:

Cycle	Date	Everolimus	Physician	Consultant
C# __				
C# __				
C# __				
C# __				
C# __				
C# __				
C# __				

Important Notes:

Reported grade 3/4 toxicities: None Hematological Non-Hematological

If yes; Did it indicate hospitalization? Yes No

Did it indicate chemo-delay for ≥ 7 days? Yes No

Did it indicate dose reduction? Yes No

Did it indicate G-CSF support? Yes No