



مركز الكويت لمكافحة السرطان
Kuwait Cancer Control Center

GEMcitabine / CARBOplatin (3 weeks)



Ministry of Health



Name:

File #:

Ht (cm):

Nationality:

Civil ID:

Wt (Kg):

Gender/Age:

DOB:

BSA (m²):

Indication(s): Multiple malignancies, specify: _____

Central line: Available NA

Allergies: NKA Yes, specify; _____

Parameters: Initiate treatment only if ANC ≥ 1000; HB ≥ 80; Plt ≥ 100,000; CrCl > 45 ml/min.

Pre-treatment Medications: (30-60 min before starting treatment)

Ondansetron 8 mg PO/IV

Dexamethasone 10 mg PO/IV

Standard Protocol:

DRUG	DOSE	ADMINISTRATION	DAYS
GEMcitabine	1000 mg/m ²	IV in 250 mL NS over 30 min.	D1, 8
CARBOplatin	AUC 5	IV in 250 mL D5W over 30 min.	D1

To be repeated every 3 weeks for 6 cycles.

Special instructions: Calvert method was used to calculate CARBOplatin dose based on serum creatinine. In case of AUC 5, the maximum CARBOplatin dose is 750 mg.

Treatment Description:

Cycle	Day	Date	GEMcitabine	CARBOplatin	Physician	Consultant
C# __	D1					
	D8			XXXXXXXX		
C# __	D1					
	D8			XXXXXXXX		
C# __	D1					
	D8			XXXXXXXX		

Important Notes:

Reported grade 3/4 toxicities: None Hematological Non-Hematological

If yes; Did it indicate hospitalization? Yes No

Did it indicate chemo-delay for ≥ 7 days? Yes No

Did it indicate dose reduction? Yes No

Did it indicate G-CSF support? Yes No