



مركز الكويت لمكافحة السرطان
Kuwait Cancer Control Center

PACLItaxel / CARBOplatin (every 3 weeks)



Ministry of Health



Name:

File #:

Ht (cm):

Nationality:

Civil ID:

Wt (Kg):

Gender/Age:

DOB:

BSA (m²):

Indication(s): Multiple malignancies, specify: _____

Central line: Available NA

Allergies: NKA Yes, specify; _____

Parameters: Initiate treatment only if ANC ≥ 1000; HB ≥ 80; Plt ≥ 100,000; CrCl > 45 ml/min.

Pre-treatment Medications: (30-60 min before starting treatment)

Ondansetron 8 mg PO/IV

Dexamethasone 10 mg PO/IV

Standard Protocol:

DRUG	DOSE	ADMINISTRATION	DAYS
PACLItaxel	175 mg/m ²	IV in 500 mL D5W glass bottle over 3 hrs.	D1
CARBOplatin	AUC 5-6	IV in 250 mL D5W over 30 min.	D1

Induction: to be repeated every 3 weeks for 2 cycles.
 Adjuvant: to be repeated every 3 weeks for 4 cycles.
 Palliative: to be repeated every 3 weeks for 6 cycles (may extend to 9 cycles).

Special instructions: Calvert method was used to calculate CARBOplatin dose based on serum creatinine.
 In case of AUC 5, the maximum CARBOplatin dose is 750 mg.
 In case of AUC 6, the maximum CARBOplatin dose is 900 mg.

Treatment Description:

Cycle	Date	PACLItaxel	CARBOplatin	Physician	Consultant
C# __					
C# __					
C# __					
C# __					
C# __					
C# __					

Important Notes:

Reported grade 3/4 toxicities: None Hematological Non-Hematological

If yes; Did it indicate hospitalization? Yes No

Did it indicate chemo-delay for ≥ 7 days? Yes No

Did it indicate dose reduction? Yes No

Did it indicate G-CSF support? Yes No