PACLItaxel / CISplatin (every 3 weeks)







Printed: 13/May/2020

Name: Nationality: Gender/Age:			File #: Civil ID: DOB:		Ht (cm): Wt (Kg): BSA (m²):
Indication(s): Multiple malignancies, specify:					
Parameters: Initiate treatment only if ANC ≥ 1000; HB ≥ 80; Plt ≥ 100,000; CrCl > 45 ml/min.					
Pre-treatment Medications: (30-60 min before starting treatment) Akynzeo 1 Capsule PO (300 mg NETUpitant/0.5 mg PALONOsetron) on Day 1 Dexamethasone 12 mg PO/IV Chlorphenamine 10 mg PO/IV					
Standard Protocol:					
DRUG DOSE		SE .	ADMINISTRATION		DAYS
PACLIta	CLItaxel 135 mg/m²		IV in 500 mL D5W glass bottle over 3 hrs.		D1
CISplatin 75		ng/m²	IV in 1000 mL NS over 2 hrs.		D1
To be repeated every 3 weeks for 6 cycles.					
Treatment Description: Pre-hydration: 1 Liter NS + KCl 20 mEq + MgSO ₄ 8 mEq IV over 1 hrs.					
Cycle	Date	PACLItaxel	CISplatin	Physician	Consultant
C#					
Post-hydration: 1 Liter NS IV over 1 hrs followed by 150 mL Mannitol 20% IV bolus.					
Pre-hydration: 1 Liter NS + KCl 20 mEq + MgSO₄ 8 mEq IV over 1 hrs.					
Cycle	Date	PACLItaxel	CISplatin	Physician	Consultant
C#					
Post-hydration: 1 Liter NS IV over 1 hrs followed by 150 mL Mannitol 20% IV bolus.					
Important Notes: Reported grade 3/4 toxicities: □ None □ Hematological □ Non-Hematological If yes; Did it indicate hospitalization? □ Yes □ No Did it indicate chemo-delay for ≥ 7 days? □ Yes □ No Did it indicate dose reduction? □ Yes □ No					

Did it indicate G-CSF support?

☐ No

☐ Yes