



مركز الكويت لمكافحة السرطان
Kuwait Cancer Control Center

TOPotecan (3 weeks)



* T O P - 0 0 0 0 - 0 3 - C N - M L *



Ministry of Health

Name:

File #:

Ht (cm):

Nationality:

Civil ID:

Wt (Kg):

Gender/Age:

DOB:

BSA (m²):

Indication(s): Small Cell Lung Cancer (Subsequent Line).

Ovarian Cancer / Cervical cancer

Central line: Available NA

Allergies: NKA Yes, specify; _____

Parameters: Initiate treatment only if ANC \geq 1000; HB \geq 80; Plt \geq 100,000; CrCl $>$ 45 ml/min.

Pre-treatment Medications: (30-60 min before starting treatment)

Ondansetron 8 mg PO/IV

Dexamethasone 10 mg PO/IV

Standard Protocol:

DRUG	DOSE	ADMINISTRATION	DAYS
TOPotecan	1.5 mg/m ²	IV in 100 mL NS over 30 min.	D1, 2, 3, 4, 5
To be repeated every 3 weeks until disease progression or intolerable toxicity.			

Treatment Description:

Cycle	Day	Date	TOPotecan
C# __	D1		
	D2		
	D3		
	D4		
	D5		

Important Notes:

Reported grade 3/4 toxicities: None Hematological Non-Hematological

If yes; Did it indicate hospitalization? Yes No

Did it indicate chemo-delay for \geq 7 days? Yes No

Did it indicate dose reduction? Yes No

Did it indicate G-CSF support? Yes No

Physician (Stamp and signature)

Consultant (Stamp and signature)