



مركز الكويت لمكافحة السرطان
Kuwait Cancer Control Center

CETUXimab Concomitant with Radiotherapy



Ministry of Health

Name:

File #:

Ht (cm):

Nationality:

Civil ID:

Wt (Kg):

Gender/Age:

DOB:

BSA (m²):

Indication(s): Locally advanced head And neck cancers.

Central line: Available NA

Allergies: NKA Yes, specify; _____

Parameters: Initiate treatment only If ANC \geq 1500; HB \geq 80; Plt \geq 100,000; CrCl $>$ 45 ml/min.

Standard Protocol:

DRUG	DOSE	ADMINISTRATION	DAYS
CETUXimab	400 mg/m ² (Loading dose)	IV undiluted over 2 hr. Then, flush the IV line with 50 mL NS at end of infusion.	1st dose only
CETUXimab	250 mg/m ² (maintenance dose)	IV undiluted over 60 min. Then, flush the IV line with 50 mL NS at end of infusion.	D1

To be given on weekly basis as long as the patient on radiation therapy.

Special instructions:

- The loading dose should be given 1 week before starting radiotherapy.
- Continue CETUXimab on weekly basis as long as the patient on radiotherapy.
- For nurses: Patients are to be observed visually for the first 15 min of each cycle of cetuximab infusion. Keep the patient for 1 hr observation period after completion of the first 2 cycles. May discontinue observation period, if no infusion reaction reported for 2 consecutive doses.

Treatment Description:

Cycle	Date	CETUXimab	Physician	Consultant
W# __				
W# __				
W# __				
W# __				
W# __				
W# __				

Important Notes:

Reported grade 3/4 toxicities: None Hematological Non-Hematological

If yes; Did it indicate hospitalization? Yes No

Did it indicate chemo-delay for \geq 7 days? Yes No

Did it indicate dose reduction? Yes No

Did it indicate G-CSF support? Yes No