



مركز الكويت لمكافحة السرطان
Kuwait Cancer Control Center

Temozolomide Concomitant with Radiotherapy



Ministry of Health

Name:

File #:

Ht (cm):

Nationality:

Civil ID:

Wt (Kg):

Gender/Age:

DOB:

BSA (m²):

Indication(s): High grade glioma.

Central line: Available NA

Allergies: NKA Yes, specify; _____

Parameters: Initiate treatment only If ANC \geq 1500; HB \geq 80; Plt \geq 100,000; CrCl $>$ 45 ml/min.

Standard Protocol:

DRUG	DOSE	ADMINISTRATION
Temozolomide	75 mg/m ² PO daily	To be given on an empty stomach or at bedtime to reduce nausea and vomiting.
Start with D1 of radiotherapy and to continue on daily basis during the external beam radiation therapy (EBRT), preferably 1hr prior to radiotherapy and in A.M. on days without radiotherapy.		

Treatment Description:

Cycle	Date	Temozolomide	Physician	Consultant
W# __				
W# __				
W# __				
W# __				
W# __				
W# __				

Important Notes:

Reported grade 3/4 toxicities: None Hematological Non-Hematological

If yes; Did it indicate hospitalization? Yes No

Did it indicate chemo-delay for \geq 7 days? Yes No

Did it indicate dose reduction? Yes No

Did it indicate G-CSF support? Yes No