



مركز الكويت لمكافحة السرطان
Kuwait Cancer Control Center

Pemetrexed (Alimta®)



* P E M - 0 0 0 0 0 - 0 3 - C N - T X *



Ministry of Health

Name:

File #:

Ht (cm):

Nationality:

Civil ID:

Wt (Kg):

Gender/Age:

DOB:

BSA (m²):

Indication(s): Early/Late stages lung adenocarcinoma.

Mesothelioma.

Central line: Available NA

Allergies: NKA Yes, specify; _____

Parameters: Initiate treatment only if ANC \geq 1000; HB \geq 80; Plt \geq 100,000; CrCl > 45 ml/min.

Preparatory Medications: To be started 1 week pre- and continue for 3 weeks post-Pemetrexed

- Folic acid 5 mg PO daily

- Vitamin B12 1000mcg IM every 9 weeks

Pre-treatment Medications: (30-60 min before starting treatment)

Ondansetron 8 mg PO/IV

Dexamethasone 10 mg PO/IV

Standard Protocol:

DRUG	DOSE	ADMINISTRATION	DAYS
Pemetrexed	500 mg/m ²	IV in 100 mL NS over 10 min.	D1
To be repeated every 3 weeks until disease progression or intolerable toxicity.			

Treatment Description:

Cycle	Date	Pemetrexed	Physician	Consultant
C# __				
C# __				
C# __				
C# __				
C# __				
C# __				

Important Notes:

Reported grade 3/4 toxicities: None Hematological Non-Hematological

If yes; Did it indicate hospitalization? Yes No

Did it indicate chemo-delay for \geq 7 days? Yes No

Did it indicate dose reduction? Yes No

Did it indicate G-CSF support? Yes No