

MEDICAL ONCOLOGY DEPARTMENT

G8 questionnaire

	Items	Possible answers (score)
A	Has food intake declined over the past 3 months due to loss of appetite, digestive problems, chewing or swallowing difficulties?	0 : severe decrease in food intake
		1 : moderate decrease in food intake
		2 : no decrease in food intake
B	Weight loss during the last 3 months	0 : weight loss > 3 kg
		1 : does not know
		2 : weight loss between 1 and 3 kgs
		3 : no weight loss
C	Mobility	0 : bed or chair bound
		1 : able to get out of bed/chair but does not go out
		2 : goes out
E	Neuropsychological problems	0 : severe dementia or depression
		1 : mild dementia or depression
		2 : no psychological problems
F	Body Mass Index (BMI (weight in kg) / (height in m ²))	0 : BMI < 19
		1 : BMI = 19 to BMI < 21
		2 : BMI = 21 to BMI < 23
		3 : BMI = 23 and > 23
H	Takes more than 3 medications per day	0 : yes
		1 : no
P	In comparison with other people of the same age, how does the patient consider his/her health status?	0 : not as good
		0.5 : does not know
		1 : as good
		2 : better
	Age	0 : >85
		1 : 80-85
		2 : <80
	TOTAL SCORE	

Result interpretation

The G-8 Score is a screening tool containing 8 questions. The total G-8 score lies between 0 and 17. A higher score indicates a better health status.

It should be noted that the G-8 tool is not aimed at replacing expertise of geriatricians for the diagnosis of frailty. Rather, it should be used as a screening tool to identify patients in need for a further assessment and appropriate care.

A threshold is suggested at 14 points, meaning that a patient with a score of 14 or lower should undergo full geriatric evaluation.